

LITERATURA

1. Amador MD, Muratet F, Teyssou E, et al. Spastic paraplegia due to recessive or dominant mutations in ERLIN2 can convert to ALS. *Neurol Genet.* 2019;5(6):e374. doi:10.1212/NXG.0000000000000374.
2. Arkadir D, Noreau A, Goldman JS, et al. Pure hereditary spastic paraplegia due to a de novo mutation in the NIPA1 gene. *Epilepsia.* 2014;55(4):e36–e39. doi:10.1111/ene.12284.
3. Bürger JM, Hazan J, Martin JJ. Hereditární spastické paraplegie: Genetická heterogenita a fenotypové projevy. *J Neurol.* 2000;247(Suppl 5):34–42.
4. Du J, Hu Y, Tang B, et al. Expansion of the phenotypic spectrum of SPG6 caused by mutation in NIPA1. *Clin Neurol Neurosurgery.* 2011;113(6):480–482. doi:10.1016/j.clineuro.2011.02.011.
5. Fink J, Klobusiakova P, Balazova Z, et al. Hereditäre spastische Paraplegien. *Nervenarzt.* 2003;74(6):497–504. doi:10.1007/s00115-003-1516-3.
6. Fu J, Ma M, Li G, et al. Clinical and Genetic Features of Chinese Patients With NIPA1-Related Hereditary Spastic Paraplegia Type 6. *Front Genet.* 2022;13:859688. doi:10.3389/fgene.2022.859688.
7. Fullam T, Statland J. Upper motor neuron disorders: primary lateral sclerosis, upper motor neuron dominant amyotrophic lateral sclerosis, and hereditary spastic paraplegia. *Brain Sci.* 2021;11(6):611. doi:10.3390/brainsci11060611.
8. Goytain A, Hines RM, El-Husseini A, et al. NIPA1 (SPG6), the basis for autosomal dominant form of hereditary spastic paraplegia, encodes a functional Mg²⁺ transporter. *J Biol Chem.* 2007;282(11):8060–8068. doi:10.1074/jbc.M610314200.
9. Harding AE, Klobusiakova P, Balazova Z, et al. Classification of the hereditary ataxias and paraplegias. *Lancet.* 1983;1(8334):1151–1155.
10. Hart CG, Karimi-Abdolrezaee S, Klobusiakova P, et al. Bone morphogenetic proteins: New insights into their roles and mechanisms in CNS development, pathology and repair. *Exp Neurol.* 2020;334:113455. doi:10.1016/j.expneurol.2020.113455.
11. Klebe S, Lacour A, Durr A, et al. NIPA1 (SPG6) mutations are a rare cause of autosomal dominant spastic paraplegia in Europe. *Neurogenetics.* 2007;8:155–157. doi:10.1007/s10048-006-0074-9.
12. Kume K, Kamada M, Shimatani Y, et al. Novel monoallelic variant in ERLIN2 causes spastic paraplegia converted to amyotrophic lateral sclerosis. *J Neurol Sci.* 2021;430:119984. doi:10.1016/j.jns.2021.119984.
13. Martinez-Lage M, Molina-Porcel L, Falcone D, et al. TDP-43 Pathology in a Case of Hereditary Spastic Paraplegia with a NIPA1/SPG6 Mutation. *Acta Neuropathol.* 2012;124(3):285–291. doi:10.1007/s00401-012-0947-y.
14. Radziwonik-Frączyk W, Elert-Dobkowska E, Kubalska J, et al. Co-occurrence of CAPN3 homozygous mutation and CCTG expansion in the CNBP gene in a patient with muscular dystrophy. *Adv Psychiatry Neurol.* 2024;33(2):109–114. doi:10.5114/ppn.2024.141382.
15. Ruano L, Melo C, Silva MC, et al. The global epidemiology of hereditary ataxia and spastic paraplegia: A systematic review of prevalence studies. *Neuroepidemiology.* 2014;42:174–183.
16. Schymick J, Talbot K, Traynor B. Genetics of sporadic amyotrophic lateral sclerosis. *Hum Mol Genet.* 2007;16(Suppl 2):R233–R242. doi:10.1093/HMG/DDM215.
17. Tanti M, Cairns D, Mirza N, et al. Is NIPA1-associated hereditary spastic paraplegia always 'pure'? Further evidence of motor neurone disease and epilepsy as rare manifestations. *Neurogenetics.* 2020;21:305–308. doi:10.1007/s10048-020-00619-0.
18. Teyssou E, Chartier L, Amador MD, et al. Novel UBQLN2 mutations linked to amyotrophic lateral sclerosis and atypical hereditary spastic paraplegia phenotype through defective HSP70-mediated proteolysis. *Neurobiol Aging.* 2017;58:239.e11–239.e20. doi:10.1016/j.neurobiolaging.2017.06.018.
19. Theuriat J, Pegat A, Leblanc P, et al. Phenocconversion from spastic paraplegia to ALS/FTD associated with CY-P7B1 compound heterozygous mutations. *Genes (Basel).* 2021;12(12):1876. doi:10.3390/genes12121876.
20. Tsang HT, Edwards TL, Wang X, et al. The hereditary spastic paraplegia proteins NIPA1, spastin, and spartin are inhibitors of mammalian BMP signalling. *Hum Mol Genet.* 2009;18(20):3805–3821. doi:10.1093/hmg/ddp324.
21. Uhrová Mészárosová A, Mazanec R, Seeman P. Hereditární spastické paraparézy: klinické a genetické aspekty. *Neurol. praxi.* 2016;17(6):373–376.
22. Zhao J, Matthies DS, Botzolakos EJ, et al. Hereditary spastic paraplegia-associated mutations in the NIPA1 gene and its Caenorhabditis elegans homolog trigger neural degeneration in vitro and in vivo through a gain-of-function mechanism. *J Neurosci.* 2008;28(51):13938–13951. doi:10.1523/JNEUROSCI.4668-08.2008.
23. Zumrová A, Mazanec R, Vyhňálek M, et al. Concomitancy of mutation in FRDA gene and FMR1 premutation in 58-year-old woman. *Neuro Endocrinol Lett.* 2005;26(1):71–74.



23.

symposium praktické neurologie

11.–12. 6. 2026
BRNO



MÍSTO KONÁNÍ:
Hotel International Brno
Husova 200/16, 602 00 Brno

